

## ORIGINAL ARTICLE

# The effectiveness of *Nicotiana tabacum* leaf extract as antibacterial agent against *Aggregatibacter actinomycetemcomitans*: an experimental study

Theresa Christy Vinsensius<sup>1</sup>  
Wiena Widyastuti<sup>2\*</sup>

<sup>1</sup>Undergraduate Program, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia

<sup>2</sup>Department of Conservation, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia

\*Correspondence:  
[wiena@trisakti.ac.id](mailto:wiena@trisakti.ac.id)

Received: 07 February 2025

Revised: 12 June 2025

Accepted: 08 July 2025

Published: 31 July 2025

DOI: [10.24198/pjd.vol37no2.61637](https://doi.org/10.24198/pjd.vol37no2.61637)

p-ISSN [1979-0201](#)

e-ISSN [2549-6212](#)

## Citation:

Vinsensius, TC; Widyastuti, W; The effectiveness of *Nicotiana tabacum* leaf extract as antibacterial agent against *Aggregatibacter actinomycetemcomitans*: an in vitro study. Padj J Dent, July. 2025; 37(2): 179-186.

## ABSTRACT

**Introduction:** Dental caries is often associated with bacterial pathogens such as *Aggregatibacter actinomycetemcomitans*, which lower the oral pH and contribute to tooth damage. Calcium hydroxide (Ca(OH)<sub>2</sub>) is commonly used in vital pulp therapy due to its antibacterial properties; however, it may cause adverse effects, including pulp necrosis. As a natural alternative, tobacco leaf extract (*Nicotiana tabacum*) extract has shown potential due to its content of antibacterial compounds such as flavonoids and terpenoids. This study aims to analyse the antibacterial effect of *Nicotiana tabacum* leaf as an antibacterial agent against *A. actinomycetemcomitans*. **Methods:** An in-vitro experimental laboratory research with a post-test-only control group design. The antibacterial activity was tested using the plate count method on samples from seven different groups. The test was conducted using the microdilution method in 96-well plates, followed by the plate count method for bacterial enumeration. The test samples included *Nicotiana tabacum* leaf extract at concentrations of 3.125%, 6.25%, 12.5%, 25%, and 50%, with calcium hydroxide serving as the positive control and distilled water as the negative control. The data were analysed using one-way ANOVA followed by Tukey' HSD post hoc test. **Results:** The minimum inhibitory concentration (MIC) of the extract against *Aggregatibacter actinomycetemcomitans* of 12.5% while a concentration of 25% represented the minimum bactericidal concentration (MBC). The 25% concentration exhibited antibacterial activity comparable to that of calcium hydroxide. The Tukey HSD post hoc test revealed significant differences between the negative control and the *Nicotiana tabacum* leaf extract at concentrations of 6.25% (p=0.025), 12.5% (p=0.001), 25% (p=0.001), and 50% (p=0.001). This study aims to analyse the antibacterial effect of *Nicotiana tabacum* leaf as an antibacterial agent against *A. actinomycetemcomitans*. **Conclusion:** *Nicotiana tabacum* leaf extract exhibits antibacterial activity against *Aggregatibacter actinomycetemcomitans*, with an MIC of 12.5% and an MBC of 25%.

## KEYWORDS

*Nicotiana tabacum* leaf extract; antibacterial effects; *Aggregatibacter actinomycetemcomitans*; calcium hydroxide

## INTRODUCTION

Dental caries is an infectious disease caused by bacterial interactions and is commonly found in humans.<sup>1</sup> In Indonesia, dental caries ranks as the most prevalent chronic disease among the population.<sup>2</sup> According to the 2023 SKI survey, 56.9% of Indonesians aged over three years experience dental and oral health issues, with a national caries prevalence reaching 88.80%.<sup>3</sup> One of the biofilm-forming microorganisms in the oral cavity is *Aggregatibacter actinomycetemcomitans*, a gram-negative bacterium that thrives under anaerobic

conditions.<sup>4</sup> The presence of *A. actinomycetemcomitans* can lower the pH of the oral cavity below 5.5, initiating the demineralization of hydroxyapatite crystals in enamel and proteolytic degradation of hard dental tissue structures.<sup>5</sup> Studies have shown that *A. actinomycetemcomitans* is found in 33.9% of deep carious lesions and significantly contributes to reversible pulpitis.<sup>6</sup>

Untreated tooth decay can lead to pain.<sup>7</sup> One way to maintain pulp vitality is by maintaining oral hygiene and treating teeth affected by deep caries.<sup>8</sup> A common treatment for caries involving the vital pulp is pulp capping using calcium hydroxide (Ca(OH)<sub>2</sub>), which has long been considered as the gold standard for this therapy.<sup>9</sup> Ca(OH)<sub>2</sub> is a strong base with a pH of 12.5 and is known for its strong antibacterial properties as well as its ability to stimulate the formation of dentin bridges in areas of exposed pulp.<sup>10</sup> However, due to its high pH, Ca(OH)<sub>2</sub> can cause superficial necrosis of the pulp surface, which may lead to re-opening of the dentin bridge.<sup>11</sup> In addition, the application of Ca(OH)<sub>2</sub> at certain concentrations may also result in fibroblast cell death, leading to pulp tissue damage.<sup>12</sup>

Currently, natural materials are widely used as alternative treatments to reduce the side effects of chemical substances in healthcare.<sup>13</sup> One natural material commonly used in Indonesia is the tobacco plant (*Nicotiana tabacum*). Tobacco leaves have long been used by the kretek cigarette industry as the main ingredient, especially to create the distinctive aroma of kretek cigarettes.<sup>14</sup> Research has shown that in addition to their role in the tobacco industry, *Nicotiana tabacum* leaf extracts contain various active compounds with potential antibacterial and antifungal properties, which may help overcome some limitations of Ca(OH)<sub>2</sub>, as the gold standard in vital pulp treatment. However, it is important to note that *Nicotiana tabacum* also contains toxic compounds, such as nicotine and other alkaloids, which may pose cytotoxic effects; therefore, its safety and dosage must be carefully evaluated prior to clinical application.<sup>15</sup>

Previous studies have tested the antibacterial activity of *Nicotiana tabacum* leaf extract against *Streptococcus mutans*, *Porphyromonas gingivalis*, and *Candida albicans* using the disc diffusion method. The results showed that a 60% concentration of *Nicotiana tabacum* leaf extract effectively inhibited these oral pathogens, with inhibition zones increased proportionally with higher extract concentrations.<sup>16</sup>

The novelty of this research lies in the application of *Nicotiana tabacum* for a bacterial target that has not been previously studied in this context, using a methodological approach distinct from previous studies. This study aims to analyse the antibacterial effect of *Nicotiana tabacum* leaf as an antibacterial agent against *A. actinomycetemcomitans*.

## METHODS

This study was an in-vitro experimental laboratory research with a post-test-only control group design. The research was conducted at the MiCORE Laboratory, Faculty of Dentistry, Universitas Trisakti, from September to November 2024. The procedure began with the preparation of leaf extract samples. Virginia tobacco (*Nicotiana tabacum*) leaves aged 2 to 3 months characterized by a healthy, green, unblemished appearance, were selected and processed into powder form. The powdered leaves were then macerated and dissolved using 96% ethanol at a ratio of 1:5 (tobacco leaf powder to ethanol). The resulting mixture was filtered using flannel cloth, and the ethanol solvent was removed using a rotary evaporator.

The extract was diluted into five concentrations (50%, 25%, 12.5%, 6.25%, and 3.125%) using sterile distilled water. In this study, a 10-0% concentration of *Nicotiana tabacum* leaf extract was not tested, as previous research had already proven that a 60% concentration is effective in eliminating oral cavity microbes, including *S. mutans*, *P. gingivalis*, and *Candida albicans*. The dilution process was calculated using the formula  $V_1 \times M_1 = V_2 \times M_2$ , where  $V_1$  represents the volume

of the pure solution (in microliters), M1 is the concentration of the stock solution (in percent), V2 is the total volume of the final solution (in microliters), and M2 is the desired concentration of the final solution (in percent).<sup>17</sup>

Bacterial media preparation is conducted by weighing 7.4 grams of Brain Heart Infusion Broth powder (Merck KGaA, Darmstadt, Germany) and 3 grams of bacteriological agar powder using a digital scale. These were then dissolved in 200 mL of sterile distilled water in an Erlenmeyer flask. The mixture was stirred with a glass rod until homogeneous, forming Brain Heart Infusion Agar (Merck KGaA, Darmstadt, Germany). The flask was covered with aluminum foil and sterilized in an autoclave at 121°C and 1 atm pressure for 15 minutes. After autoclaving, the BHI-A medium was poured into seven sterile Petri dishes to a thickness of approximately 4 mm and left to solidify.<sup>17</sup>

The bacterial sample used was *A. actinomycetemcomitans* ATCC 29522. The bacteria were suspended in 10 mL of BHI-B, vortexed, and incubated. A microplate reader was used to measure the turbidity and determine the bacterial colonies, with the turbidity standard following McFarland 0.5 or  $1.5 \times 10^8$  CFU/mL. The antibacterial activity test was conducted using the microdilution method in 96-well plates, followed by the plate count method for bacterial enumeration.<sup>17</sup>

The preparation of the positive control began by mixing the base and catalyst components of calcium hydroxide ( $\text{Ca}(\text{OH})_2$ ) from Dentsply using a cement spatula on a paper pad until evenly blended. Then, 0.28 grams of the  $\text{Ca}(\text{OH})_2$  mixture was placed into a microcentrifuge tube, followed by the addition of 0.7 mL of distilled water, and vortexed for 15 seconds. A micropipette was used to dispense the resulting solution into 96-well plates with four replicates, and the mixture was then combined with the bacterial suspension.<sup>17</sup>


The antibacterial test against *A. actinomycetemcomitans* was performed using a dilution method involving seven test groups: *Nicotiana tabacum* leaf extract at concentrations of 3.125%, 6.25%, 12.5%, 25%, and 50%, distilled water as a negative control, and  $\text{Ca}(\text{OH})_2$  as a positive control. A micropipette is used to dispense 100  $\mu\text{L}$  of *A. actinomycetemcomitans* bacterial suspension and dispense it into each well of the 96-well plates, followed by the addition of 100  $\mu\text{L}$  of each test solution. The mixture was then incubated for 24 hours at 37°C under anaerobic conditions.<sup>17</sup>

After incubation, each mixture of bacterial suspension and test solution was diluted 1000 times using a micropipette. Then, 5  $\mu\text{L}$  of the diluted solution was streaked onto petri dishes containing BHI-A medium and incubated for 24 hours at 37°C under anaerobic conditions. Bacterial colony counts were conducted to determine the minimum inhibitory concentration (MIC). The BHI-A media was further incubated for another 24 hours to determine the minimum bactericidal concentration (MBC). Bacterial colonies were counted four times on each petri dish using the plate count method.<sup>17</sup>

The data were then analysed by applying the Shapiro-Wilk test to evaluate the normality. If the data were normally distributed ( $p > 0.05$ ), a parametric test using one-way analysis of variance (ANOVA) was performed. When significant differences were detected ( $p < 0.05$ ), the Tukey HSD (honestly significant difference) post hoc test was used to identify specific groups differences.

## RESULTS

The results of antibacterial tests for all groups treated with *Nicotiana tabacum* leaf extract are presented in the following figure.



**Figure 1.** Antibacterial test results of *Nicotiana tabacum* leaf extract against *A. actinomycetemcomitans* using the plate count method. (A) *Nicotiana tabacum* leaf extract 50%; (B) *Nicotiana tabacum* leaf extract 25%; (C) *Nicotiana tabacum* leaf extract 12.5%; (D) *Nicotiana tabacum* leaf extract 6.25%; (E) *Nicotiana tabacum* leaf extract 3.125%; (F) Ca(OH)<sub>2</sub> (positive control); (G) Aquadest (negative control).

On petri dishes containing *A. actinomycetemcomitans*, no bacterial colonies were observed in the groups treated with *Nicotiana tabacum* leaf extract at concentrations of 50%, 25%, as well as in the Ca(OH)<sub>2</sub> positive control group. Treatment with *Nicotiana tabacum* leaf extract at a concentration of 12.5% resulted in an average colony count of  $(239 \pm 215) \times 10^5$  CFU/mL. At a 6.25%, the average colony count was  $(423 \pm 66.42) \times 10^5$  CFU/mL, and at a 3.125%, the average count increased to  $(642 \pm 132.12) \times 10^5$  CFU/mL. The negative control group (distilled water) exhibited the highest average colony count, at  $(716 \pm 43.99) \times 10^5$  CFU/mL (Table 1). *Nicotiana tabacum* leaf extracts contain active compounds with antibacterial effects against *A. actinomycetemcomitans* as presented in the Table 1 below.

**Table 1. Average colony count of *Aggregatibacter actinomycetemcomitans***

Concentration Group	Average number of colonies ± SD (CFU/mL)
A <i>Nicotiana tabacum</i> leaf extract 50%	$(0.00 \pm 0) \times 10^5$
B <i>Nicotiana tabacum</i> leaf extract 25%	$(0.00 \pm 0) \times 10^5$
C <i>Nicotiana tabacum</i> leaf extract 12,5%	$(239 \pm 215) \times 10^5$
D <i>Nicotiana tabacum</i> leaf extract 6,25%	$(423 \pm 66.42) \times 10^5$
E <i>Nicotiana tabacum</i> leaf extract 3,125%	$(642 \pm 132.12) \times 10^5$
F Ca(OH) <sub>2</sub>	$(0.00 \pm 0) \times 10^5$
G Aquadest	$(716 \pm 43.99) \times 10^5$

The normality test showed that the data were normally distributed ( $p > 0.05$ ), allowing for a one-way ANOVA test. Table 2 indicates that the p-value is extremely small ( $p < 0.001$ ), so it's not displayed numerically by the statistical software used. This suggests that the data at these concentrations show significant difference from the negative control.

**Table 2. Tests of normality results using Shapiro-Wilk**

	K-	3.125%	6.25%	12.5%	25%	50%	K+
p value	0.220	0.057	0.533	0.505	<0.001	<0.001	<0.001

The one-way ANOVA revealed a significant difference among the test groups ( $p < 0.05$ ). Therefore, the analysis was followed by the Tukey HSD post hoc test. The results indicated significant differences between the negative control and the *Nicotiana tabacum* leaf extract at concentrations of 6.25%, 12.5%, 25%, and

50%. However, no significant difference was observed at a concentration of 3.125% (Table 3).

**Table 3. Antibacterial test results of *Nicotiana tabacum* leaf extract compared to negative and positive controls using Tukey HSD post hoc test**

	3.125%	6.25%	12.5%	25%	50%
K (-)	0.967	0.025*	0.001*	0.001*	0.001*
K (+)	0.001*	0.001*	0.095	1	1

\* = significant difference with a p-value < 0.05

Based on these results, the minimum inhibitory concentration (MIC) of *Nicotiana tabacum* leaf extract against *Aggregatibacter actinomycetemcomitans* was determined to be 12.5% because it began to inhibit bacterial growth. The minimum bactericidal concentration (MBC) was determined to be 25%, since no bacterial colonies were observed at this concentration and higher.

## DISCUSSION

In this study, *Nicotiana tabacum* leaves were macerated with 96% ethanol to obtain the extract. The maceration technique is a preferred method, as it minimizes the risk of damaging the active compounds present in *Nicotiana tabacum* leaves during extraction process.<sup>18</sup> The use of 96% ethanol as a solvent is considered effective due its low toxicity and high solubility.<sup>19</sup> According to the literature, ethanol-based solvents can optimize the extraction of flavonoids and phenolic compounds found in plants.<sup>20</sup>

The findings of this study in Table 1 showed that *Nicotiana tabacum* leaf extract has antibacterial activity against *A. actinomycetemcomitans*. At a concentration of 12.5%, bacterial growth was still observed with an average colony count of  $(239 \pm 215) \times 10^5$  CFU/mL. In contrast, no bacterial growth was detected at concentrations of 25% and 50% concentrations. The MIC value was defined as the lowest concentration of *Nicotiana tabacum* leaf extract that inhibited visible bacterial growth, while the MBC referred to the lowest concentration that completely eliminated bacterial colonies.<sup>21</sup> Based on these criteria, the 12.5% concentration was identified as the MIC, and the 25% concentration was determined to be as the MBC.

The antibacterial test was conducted on *A. actinomycetemcomitans*, one of the bacteria commonly associated with deep caries. Along with other deep pathogens, *A. actinomycetemcomitans* can contribute to pulp irritation.<sup>6</sup> Calcium hydroxide ( $\text{Ca}(\text{OH})_2$ ) was used as the positive control in this study due its potent antibacterial properties, making it known as the gold standard for vital pulp treatment materials. In vital pulp treatment, particularly pulp capping,  $\text{Ca}(\text{OH})_2$  helps protect the pulp complex and maintain pulp vitality.<sup>9</sup> Distilled water, which has neutral properties and does not promote bacterial growth, was used as the negative control.<sup>22</sup> Antibacterial testing performed to determine the MIC and MBC values was conducted using the microdilution method, followed by the plate count method. The plate count method for bacterial colony enumeration is relatively simple and does not require complex equipment, making it accessible for laboratory use.<sup>23</sup>

The results indicated significant differences between the negative control and the *Nicotiana tabacum* leaf extract at concentrations of 6.25%, 12.5%, 25%, and 50% in Table 3, possibly because *A. actinomycetemcomitans* is a gram-negative bacterium with a cell wall containing high lipid content. In this study, it is suggested from previous study that the lipid compounds in the cell wall of *A. actinomycetemcomitans* were disrupted upon contact with the extract, causing bacterial cell walls to form pores that facilitated the penetration of active compounds from the *Nicotiana tabacum* leaf extract.<sup>24</sup> The 25% concentration demonstrated antibacterial effectiveness similar to that of  $\text{Ca}(\text{OH})_2$ , used as the



positive control. Overall, the findings indicate that the antibacterial activity of *Nicotiana tabacum* against *A. actinomycetemcomitans* increases with higher extract concentrations.

These results are in line with previous studies that evaluated the antibacterial effects of *Nicotiana tabacum* leaf extract against *Streptococcus mutans*, *Porphyromonas gingivalis*, and *Candida albicans*. Those studies, which used the well diffusion method, showed that the antibacterial activity became stronger with higher concentration -similar to the findings observed in this study.<sup>25</sup> However, other research examining *N. tabacum* extract against *Staphylococcus aureus* reported different results. In that study, the antibacterial effect did not consistently increase with higher concentrations. This inconsistency may be attributed to the formation of larger extract particles at higher concentrations, which could hinder their ability to penetrate the bacterial cell wall. In this case of *S. aureus*, this may be due to limited transport through membrane proteins.<sup>26</sup>

The antibacterial effects of *Nicotiana tabacum* leaf extract against *A. actinomycetemcomitans* may be attributed to the presence of active compounds such as flavonoids, saponins, steroids, and terpenoids. Each of these compounds has its own unique mechanism for inhibiting bacterial growth. Flavonoids, a type of polyphenol, inhibit bacterial proliferation by disrupting energy metabolism, damaging the cell membrane, and reducing nucleic acid synthesis.<sup>25</sup> Saponins and steroids have been reported to possess anti-genotoxic, anti-mutagenic, and anti-inflammatory properties. The antibacterial mechanism of terpenoids in *Nicotiana tabacum* is believed to involve the inhibition of the bacterial enzymes responsible for cell wall synthesis.<sup>26</sup>

The average total colony count of *A. actinomycetemcomitans* showed a relatively high standard deviation, which may be attributed to several limitations in the study. First, the experimental procedures were highly sensitive to technical variations, such as inconsistencies in bacterial streaking, pipetting errors, or potential cross-contamination, all of which could affect the accuracy and reproducibility of the results. Additionally, environmental factors such as pH, temperature, and media composition in the test conditions may not fully replicate clinically or from real-world settings, potentially affecting the observed antibacterial activity of the tested compounds.

These results indicate that *Nicotiana tabacum* leaf extract has the potential to serve as a promising and innovative alternative for vital pulp therapy. Its antibacterial activity against *A. actinomycetemcomitans* was similar to that of calcium hydroxide (Ca(OH)<sub>2</sub>). With further refinement and evaluation of its safety profile, *Nicotiana tabacum* may offer a natural, plant-based option for use in vital pulp treatment.<sup>27</sup>

To further explore the potential use of *Nicotiana tabacum* leaf extract as an alternative for vital pulp therapy, additional studies are recommended. These should include the use of different extraction methods to obtain a more optimal extract and its efficacy, in vitro toxicity and solubility assessments and investigations into its potential synergistic effects when combined with other biomaterials.

One limitation of this study is the use of crude extracts without isolating specific active compounds, which may result in unpredictable antibacterial effects. To address this, future studies should explore alternative extraction methods to obtain a more optimal (refined and standardized) *Nicotiana tabacum* leaf extract. Additionally further testing is recommended to within the concentration range of 12.5% to 25% to determine a more accurate minimum inhibitory concentration (MIC) value.

## CONCLUSION

*Nicotiana tabacum* leaf extract exhibits antibacterial effects against *A. actinomycetemcomitans*, with a minimum inhibitory concentration (MIC) of 12.5% and a minimum bactericidal concentration (MBC) of 25%. The 25% *Nicotiana tabacum* leaf extract demonstrated antibacterial effects equivalent to those of Ca(OH)<sub>2</sub>. The implications of this research support the potential for further investigation into plant-based antibacterial agents as alternatives to chemical-based such as Ca(OH)<sub>2</sub> for use in pulp therapy.

**Acknowledgement:** The author would like to thank the professionals who contributed to the research and preparation of this manuscript.

**Author Contributions:** Conceptualization, TCV and WW; methodology, TCV; software, TCV; validation, TCV and WW; formal analysis, TCV; investigation, TCV; resources, TCV; data curation, TCV; writing original draft preparation, TCV; writing review and editing, TCV; visualization, TCV; supervision, TCV; project administration, TCV; funding acquisition, WW. All authors have read and agreed to the published version of the manuscript.", please turn to the CRediT taxonomy for the term explanation.

**Funding:** This research received no funding.

**Institutional Review Board Statement:** Ethical review and approval were waived for this study, as it did not involve human participants or animal subjects.

**Informed Consent Statements:** Not applicable. This study did not involve human participants.

**Data Availability Statement:** Data supporting the findings of this study are available from the corresponding author upon reasonable request.

**Conflicts of Interest:** The authors declare no conflict of interest or potential commercial background related to this research.

## REFERENCES

1. Santín GR, Salgado AV, Bastida NM, de la Rosa Gómez I, Benítez JG, Zerón HM. Salivary immunoglobulin gene expression in patients with caries. *Maced J Med Sci*. 2017;5(2):236. <https://doi.org/10.3889/oamjms.2017.028>
2. Andayasari L, Mubasyiroh R, Nurlinawati I, Sufiawati I. Association between tobacco smoking and dental caries in the Indonesian population: Results of a national study in 2018. *J Prev Med Public Health*. 2023;56(4):357-67. <https://doi.org/10.3961/jpmph.22.417>
3. Ministry of Health of the Republic of Indonesia. Indonesia Health Survey 2023. Jakarta: MOH RI; 2023. pp.317-39.
4. Kozlovsky A, Wolff A, Saminsky M, Mazor Y, Venezia E, Bar-Ness Greenstein R. Effect of *Aggregatibacter actinomycetemcomitans* from aggressive periodontitis patients on *Streptococcus mutans*. *Oral Dis*. 2015;21(8):955-61. <https://doi.org/10.1111/odi.12362>
5. Dipalma G, Inchingolo AD, Inchingolo F, Charitos IA, Di Cosola M, Cazzolla AP. Focus on the cariogenic process: microbial and biochemical interactions with teeth and oral environment. *J Biol Regul Homeost Agents*. 2021;35(2):747-54. <https://doi.org/10.23812/20-747-a>
6. Radman IK, Djeri A, Arbutina A, Milašin J. Microbiological findings in deep caries lesions. *Stomatol Glas Srb*. 2019;63(1):7-14. <https://doi.org/10.1515/sdj-2016-0001>
7. Heng CC. Tooth decay is the most prevalent disease. *J Dent Res*. 2016;95(3):1-2. <https://doi.org/10.1177/0022034516632541>
8. Grigaluskienė R, Slabšinskienė E, Vasiliauskiene I. Biological approach of dental caries management. *Stomatologija*. 2015;17(4):107-12.
9. Poggio C, Beltrami R, Colombo M, Ceci M, Dagna A, Chiesa M. In vitro antibacterial activity of different pulp capping materials. *J Clin Exp Dent*. 2015;7(5):584. <https://doi.org/10.4317/jced.52401>
10. Widjiastuti I, Mudjiono M, Sudiartha NC, Samuel GA, Mega NP. Dosage effectiveness of the combination calcium hydroxide-propolis as an inhibitor for *Aggregatibacter actinomycetemcomitans* biofilm. *Indian J Forensic Med Toxicol*. 2021;15(2):2597-605. <https://doi.org/10.37506/ijfmt.v15i2.14764>
11. Ba-Hattab R, Al-Jamie M, Aldreib H, Alessa L, Alonazi M. Calcium hydroxide in endodontics: an overview. *Open J Stomatol*. 2016;6(12):274-89. <https://doi.org/10.4236/ojst.2016.612033>
12. Vincentia Adya Paramitta, Tetiana Haniastuti, Susilowati Susilowati. The Effect of Calcium Hydroxide on Fibroblast Cells Viability. *Ind J Dent Res*. 2015;1(2). <https://doi.org/10.22146/theindidentres.9996>
13. Ningtyas EA, Santoso O, Sadhana U, Sunarintyas S. Role of combination casein and lactoferrin bovine<sup>TM</sup> colostrum as a pulp capping on macrophage expression in male wistar rats. *ODONTO*. 2021;8(2):156-64. <https://doi.org/10.30659/odj.8.2.156-164>
14. P Jin, et al. E-cigarettes in ten Southeast Asian countries. *Tob Induc Dis*. 2017;15(1):1-10. [https://doi.org/10.1016/S2414-6447\(19\)30097-1](https://doi.org/10.1016/S2414-6447(19)30097-1)
15. Al-Lahham S, Sbieh R, Jaradat N, Almasri M, Mosa A, Hamayel A, Hammada F. Antioxidant, antimicrobial and cytotoxic properties of four different extracts derived from the roots of *Nicotiana tabacum* L. *Eur J Integr Med*. 2020;35:101077. <https://doi.org/10.1016/j.eujim.2019.101039>
16. Suryadi H, Sari R, Suryanto E, et al. Effect of gamma irradiation on the caffeoylquinic acid derivatives and saponin content in *Ziziphus mauritiana* leaves. *Radiat Phys Chem*. 2021;178:109216. <https://doi.org/10.1016/j.radphyschem.2020.109216>

17. Priscila K, Santosa DN. Efek potensiasi kombinasi sefadroksil dan ekstrak camellia sinensis terhadap pertumbuhan *Aggregatibacter actinomycetemcomitans* dan *Porphyromonas gingivalis*. Jurnal Kedokteran Gigi Terpadu. 2023;5(1). <https://doi.org/10.25105/jkgt.v5i1.17184>
18. Phytochemical analysis of commercial *Ziziphus mauritiana* tea: Effects of infusion time and water volume on saponin content. Malay J Analytical Sci. 2023;27(5):1089–99. doi: <https://doi.org/10.17576/mjas-2023-2705-13>
19. Viszwapriya D, Suresh S, Rajendran R, et al. In vitro and in vivo antibiofilm potential of 2,4-Di-tert-butylphenol isolated from *Eleutheria americana* Merr. Phytomedicine. 2016;23(10):1069–1076. <https://doi.org/10.1016/j.phymed.2016.06.002>
20. Alara OR, Abdurahman NH, Olalere OA, et al. Ethanolic extraction of flavonoids, phenolics and antioxidants from *Passiflora edulis* seeds: Optimization and characterization. J Food Sci Technol. 2020;57(5):1912–1923. <https://doi.org/10.1007/s11483-020-01503-4>
21. Rajapaksha RMHKK, Fernando EMN, Bandara AWMKK, Nelumdeniya NRM, Silva ARN. In-vitro anti-bacterial activity of methanol and aqueous crude extracts of *Horsfieldia iryaghedhi*. Asian Plant Res J. 2024;12(4):27–34. <https://doi.org/10.9734/aprj/2024/v12i4259>
22. Aribisala JO, Abdulsalam RA, Dweba Y, Madonsela K, Sabiu S. Identification of secondary metabolites from *Crescentia cujete* as promising antibacterial therapeutics targeting type 2A topoisomerases through molecular dynamics simulation. Comput Biol Med. 2022;145:105432. <https://doi.org/10.1016/j.compbiomed.2022.105432>
23. Kyrylenko A, Eijlander RT, Alliney G, de Bos EL, Wells-Bennik MHJ. Levels and types of microbial contaminants in different plant-based ingredients used in dairy alternatives. Int J Food Microbiol. 2023;407:110392. <https://doi.org/10.1016/j.foodcont.2023.109396>
24. Sulaiman SA, Al-Farsi M, Al-Habsi K, Al-Habsi S, Al-Saadi H, Al-Mamari H. Antibacterial activity of *Citrus aurantifolia* peel essential oil against *Aggregatibacter actinomycetemcomitans* and its synergistic effect with amoxicillin. J Appl Microbiol. 2020;128(3):723–732. <https://doi.org/10.1111/jam.14568>
25. Anumudu CK, Nwachukwu MI, Obasi CC, Nwachukwu IO, Ihenetu FC. Antimicrobial activities of extracts of tobacco leaf (*Nicotiana tabacum*) and its grounded snuff (Utaba) on *Candida albicans* and *Streptococcus pyogenes*. J Trop Dis. 2019;7(300):2. <http://dx.doi.org/10.4172/2329-891X.1000300>
26. Prommaban A, Srisuk N, Srisuk S, Chittasupho C, Sirilun S, Hemsuwimon K, Chaiyana W. Phytochemical, antimicrobial, and proximate composition of *Nicotiana tabacum* leaves extract. Phytochem Lett. 2022;50:1–9. <https://doi.org/10.1016/j.phytol.2022.01.001>
27. Jalan AL, Warhadpande MM, Dakshindas DM. A comparison of human dental pulp response to calcium hydroxide and biodentine as direct pulp-capping agents. J Conserv Dent. 2017;20(2):129–133. <http://dx.doi.org/10.4103/0972-0707.212247>